

A study on the optic glands of *Sepioteuthis lessoniana* from the Red Sea

Waheed M. Emam and Tarek A. Ghareb

Zoology Department, Faculty of Science, Ain Shams University

ABSTRACT

In the present work the optical glands of the reef squid *Lessoniana lessoniana* were found in young and adult individuals. They are paired small elongated organs that lie at the aboral end of the olfactory lobe, on a level with the optic tract. Histologically, the cells of the optical gland are large, with about 60 μm in diameter and have spherical nuclei (20 μm in diameter). Their cytoplasm is coarsely granulated. Neurosecretory granules were seen in cells of optical glands and stained with paraldehyde fuchsin. No distinct neuropil was observed in these glands and these glands are rich in blood supply. The optic glands were found to be connected with optic nerve fibres to the dorso-lateral lobes of the brain. The function of the optical glands of *Sepioteuthis lessoniana* was also discussed in this study.

Key words: Optic gland, reef squid, *Sepioteuthis lessoniana*, Cephalopods, Histology, Neurosecretion.

INTRODUCTION

The structure and function of the optic glands in dibranchiate cephalopods had been studied by many authors (Wells & Wells, 1959, 1969, 1975; Wells, 1960; Björkman, 1963; Richard, 1966, 1967 a, b, 1970; Defretin & Richard, 1967; Durchon & Richard, 1967; O'Dor & Wells 1973; Wells *et al.*, 1975; Wodinsky, 1977; Mangold, 1987; Le Gall *et al.*, 1988; Dwight, 1999).

The coral reef squid *Sepioteuthis lessoniana* (Cephalopoda: Loliginidae) represents one of the most common squids in the Red Sea. Its morphology, morphometry, age and growth for specimens from the Gulf of Suez have been studied by Emam *et al.* (2007); Emam and Ghareb (2010, 2011). Also, its spermatophoric structure was investigated by Aly (2008). However, its nervous system and optic glands were not studied till now. Therefore the present study aimed to study the structure and the proposed function of the optic glands of the coral reef squid *Sepioteuthis lessoniana* from the Gulf of Suez and compare them with those in other related cephalopods.

MATERIALS AND METHODS

Specimens of *Sepioteuthis lessoniana* were brought alive and dissected to expose the brain. The optic glands with the optic lobe and the optic stalk were immersed in Bouan fixative and then well washed in running tap water, dehydrated and embedded in paraffin. Paraffin sections (6 μm) were stained with Hematoxylin-Eosin (HE). Paraldehyde fuchsin

Waheed M. Emam and Tarek A. Ghareb

stain was also used for neurosection of optic gland cells. Also, the brain with optic glands of some specimens were prepared for Cajal silver impregnation to show nerve fibers that supply the optic glands.

RESULTS AND DISCUSSION

The optic glands are small elongated paired structure located at the posterior edge of the optic brain tract over the helium of the optic lobes and between peduncle and olfactory lobes of the brain (Figs. 1, 2 & 3). This site is similar to that in other cephalopods (Boycott & Young 1956).

Histologically cells of the optic glands are well differentiated from the cerebral tissue where they form a compact mass, whereas cerebral cells are larger, and form layers surrounding eosinophylic nerve fibers (Figs. 2 & 3). No distinct neuropil was observed in these glands. The cells of the optical gland are small in size in smaller individuals and large in mature individuals (Fig. 4), and reach about 60 μm in diameter and have spherical nuclei (20 μm in diameter). Their cytoplasm is coarsely granulated (Fig. 5). The blood vessels are shown dispersed throughout the optic glands.

Björkman (1963), Nishioka *et al.* (1966), Barber (1967), Bonichon (1967), Froesch (1974) and Mangold & Froesch (1977) reported that the optic glands in *Loligo opalescens*, *Octopus vulgaris*, *O. bimaculatus*, *Eledone moschata* and *Sepia officinalis*, consist of main or secretory cells, blood vessels, nerve cells and support cells.

The secretory cells of the optic gland in *S. lessoniana* are stained with paraldehyde fuchsin. They are the most numerous cells, and they are nearly ovoid with cytoplasmic processes and their nuclei are circular with a dense nucleoli (Figs. 6 & 7). These cells with their nuclei are larger in size in large individuals than in small ones in both sexes. Dwight (1999) showed that the organelles inside the secretory cells of the optic glands of the tropical *Sepioteuthis sepioidea* have differences in their abundance between the sexual maturity stages: at the juvenile and immature stages there are numerous mitochondriae, endoplasmic reticulum cisterns are thin and dispersed in the cytoplasm, the Golgi apparatus has a curved profile and actively secretes granules of different densities, whilst in the IV maturation stage (females) and reproductive individuals (males) secretory cells are almost depleted of mitochondriae, endoplasmic reticulum cisterns are dilated and well developed in a concentric fashion, and the Golgi complex actively secretes electron dense granules which are dispersed in the cytoplasm. Some of these are close to the lipofuchsin granules and to the cellular membrane, suggesting a possible secretory mechanism through the blood vessels. On the other hand, Dwight (1999) demonstrated the secretory activity of optic gland in *S. sepioidea* by the production of electron dense secretory granules in the sexually mature stage.

Defretin and Richard (1967) demonstrated that when *S. officinalis* is kept in a dark photoperiod the optic gland secretory cells produce three kinds of secretory granules, suggesting that they are filled with the same product in different stages of synthesis. Wells (1964) reported that the optic gland in *Octopus vulgaris* enlargement occurs both in the nucleus and secretory cell cytoplasm after cerebral surgery, while Boyle and Thorpe (1984) indicated that enlargement in optic gland of *Eledone cirrhosa* was statistically associated with female gonad maturation, and it is suspected that this process occurs normally. The cellular changes in *S. officinalis* optic gland histology was found during sexual maturation and the

A study on the optic glands of *Sepioteuthis lessoniana* from the Red Sea

main cell secretory activity was correlated with a peak of mitogenic activity in both sexes (Koueta *et al.*, 1995).

The optic glands were found to be connected with optic nerve fibres to the dorso-lateral lobes of the brain (Fig. 3).

Young (1967) mentioned that optic glands have a nervous origin and probably originated from cells of the olfactory lobes in Octopus, however Young (1977) stated that the dorso-lateral lobes are the source of fibres of these glands in Loligo and Octopus.

The role of the optic glands in *Sepioteuthis lessoniana* seems to be similar to that reported by Wells & Wells (1959, 1969); Wells (1960); for other cephalopods and is correlated with controlling of sexual development. On the other hand, Durchon & Richard (1967); O'Dor & Wells (1973); Wells *et al.* (1975); Wodinsky (1977); Mangold (1987) mentioned that these glands also regulate protein synthesis in the ovaries, stimulate gonadal mitosis, mating behavior, feeding and longevity. Wells & Wells (1975); Defretin & Richard, (1967); Richard (1966, 1967 a,b, 1970) found that the optic glands hormone is species specific, not sex specific, and glandular secretions have periodic oscillations associated with temperature and photoperiod. The glands are controlled by a reactive peptidergic innervation (Le Gall *et al.*, 1988), and their activity varies during the life cycle, associated with the multiplication of reproductive cells (Koueta *et al.*, 1995).

The optic glands of dibranchiate cephalopods are the source of a hormone that controls the state of the gonad and of its ducts. Animals can be forced into precocious maturity by cutting an inhibitory nerve to the glands from the central supra-oesophageal part of the brain (Wells & Wells, 1959, 1972). Richard (1970) has shown that fragments of the ovary from *Sepia* will carry out oogenesis-to-oocyte cell divisions in organ-culture experiments only if optic glands are present. Wells and Wells (1975) found that optic glands transplanted from one *Octopus vulgaris* into another cause enlargement of the gonads and ducts of the recipient. Enlargement occurs whether or not the gland was secreting when implanted and regardless of the sex of the donor or recipient. Also they found that optic glands derived from *Eledone moschata* or *Octopus macropus* implanted into *O. vulgaris* are as effective as glands derived from *O. vulgaris*. However, optic glands implants derived from *Sepia officinalis* or *Loligo vulgaris* appear to be ineffective.,

Wodinsky (1977) found that removal of both optic glands in cephalopods after spawning results in cessation of broodiness, resumption of feeding, increased growth, and extended life span. He stated that optic gland secretions may cause death of most cephalopods, and may control population size. Boyle and Thorpe (1984) indicated that optical gland size varies as a reproductive character together with ovary and oviducal gland size and this was not associated with other cerebral organs. Di Cosmo and Di Cristo (1998) reported the neurosecretion function of optic gland of *Octopus vulgaris*. They found that optic gland secret also gonadotropin hormone.

The fine structure of the gland suggests that it has several functions. Björkman (1963) suggested that the secretory cells had a high rate of synthesis. The presence of abundant mitochondria implies that the gland may secrete a steroid hormone (Froesch, 1979). In *S. officinalis*, however a peptidic like hormone was shown to be present in the optic glands and hemolymph (Koueta *et al.*, 1992). The lipofuchsin granules have been related to the resorption of metabolic residues inside the secretory cells (Froesch & Mangold, 1976; Mangold & Froesch, 1977; Froesch *et al.*, 1987).

Waheed M. Emam and Tarek A. Ghareb

On the other hand, the fine structure of optical gland in tropical squid *sepiotheutis sepioidea* was discussed by Dwight (1999). He found that the optic gland showed the same cellular types described in other cephalopods, with slight differences. The secretory cells in young individuals present a dendritic profile with thin cytoplasmic processes, numerous circular mitochondriae with tubular crests and abundant ribosomes. At the last stage of maturity the secretory cells contain rough endoplasmic reticulum in the form of dilated cisternae, with an active Golgi apparatus, secreting electron dense granules. Mitochondriae are scarce and lipofuchsin granules show an increase in size. Secretory cell nucleus length and organelle abundance showed oscillations associated with the life cycle.

REFERENCES

- Barber, V. (1967). A neurosecretory tissue in *Octopus*. *Nature* 213:1042-1043.
- Björkman, N. 1963. On the ultrastructure of the optic gland in *Octopus*. *J. Ultrastruct. Res.* 8:195.
- Bonichon, A. (1967). Contribution à l'étude de la neurosécrétion et de l'endocrinologie chez les Céphalopodes. I. *Octopus vulgaris*. *Vie Milieu* 18:27-263.
- Boycott, B. B. and J. Z. Young (1956). The subpedunculate body and nerve and other organs associated with the optic tracts of cephalopods. *Bertil Hanström Zoological Papers* 76-105 p.
- Defretin, R. & A. Richard 1967. Ultrastructure de la glande optique de *Sepia officinalis* L. (Mollusques; Céphalopode). Mise en évidence de la sécrétion et de son contrôle photopériodique. *C. R. Acad. Sci., Paris. D.* 265:1415-1418.
- Boyle, R.S. Thorpe (1984). Optical gland enlargement and female gonad maturation in population of octopus *Eledone cirrhosa*. *Mar. Biol.*, 79 (2):127-132.
- Di Cosmo and Di Cristo (1998). Neurosecretion control of optical gland of *Octopus vulgaris*. *J. Comp. Neurol.*, 17(398):1-12.
- Durchon, M. and A. Richard (1967). Étude en culture organotypique du rôle endocrine de la glande optique dans la maturation ovarienne chez *Sepia officinalis* L. (Mollusque: Céphalopode). *C. R. Acad. Sci., Paris. D.* 264:1497-1500.
- Dwight Arrieche (1999). Optical gland of *Sepiotheutis sepioidea*, *REV. Biol. Trop.*, 47(4):34-77.
- Emam, W.M. and Ghareb, T.A. (2010). A study on the morphology, digestive and reproductive system of male squid *Sepioteuthis lessoniana* (Cephalopoda: Loliginidae) from Abu Qir on the Mediterranean Sea. *Afr. J. Biol. Sci.*, 6(1):125-141.
- Emam, W.M. and Ghareb, T. (2011). Blood supply of Gliovascular tissue of nervous system of *Sepioteuthis lessoniana* from the Gulf of Suez. *Afr. J. Biol. Sci.*, 7(1):135-140.
- Emam, W.M.; Aly R.H.; Ibrahim, A.M. and El-Naggar, M.A. (2007). Morphology, morphometry, age and growth of the coral reef squid *Sepioteuthis lessoniana* (Cephalopoda: Loliginidae) from the Gulf of Suez. *J. Egypt. Acad. Soc. Environ. Develop., (D-Environmental Studies)* 8(3): 117–131.
- Evans, P., E. Kravitz & B. Talamo (1976). Octopamine release at two points along lobster nerve trunks. *J. Physiol.* 262:71-89.
- Froesch, D. (1979). Antigen-induced secretion in the optic gland of *Octopus vulgaris*. *Proc. R. Soc. Lond. B.* 205:379-384.

A study on the optic glands of *Sepioteuthis lessoniana* from the Red Sea

- Froesch, D., K. Mangold and W. Fritz (1978). On the turnover of exogenous ferritin in the cephalopod optic gland. A microprobe. *Experientia*, 34:116-117.
- Gervasio, A., R. Martin and A. Miralto (1971). Fine structure of synaptic contacts in the first order giant fiber system of the squid. *Z. Zellforsch*, 112:85-96.
- J. Egypt. Acad. Soc. Environ. Develop., (D-Environmental Studies), 8 (3): 117–131
- Koueta, N., E. Boucaud-Camou and A. M. Renou. (1995). Gonadotropic mitogenic activity of the optic gland of the cuttlefish *Sepia officinalis*, during sexual maturation. *J. Mar. Biol. Ass. U. K.* 75:391-404.
- Koueta, N., I. Robbins & E. Boucaud-Camou (1992). Partial characterization of a gonadotropic, mitogenic factor from the optic gland and hemolymph of the cuttlefish *Sepia officinalis*. *L. Comp. Biochem. Physiol.*, 102A:224-234.
- Le Gall, S., C. Feral, J. VanMinnen and C. Marchand (1988). Evidence for peptidergic innervation of the endocrine optic gland in *Sepia* by neurons showing FMRF amide like immunoreactivity. *Brain Res.*, 462:82-88.
- Mangold, K. and D. Froesch (1977). A reconsideration of factors associated with sexual maturation. *Symp. Zool. Soc. Lond.*, 38:541-555.
- Mangold, K. 1987. Reproduction. Chapter X. In "Cephalopod Life Cycles". Academic, New York. p. 157-200.
- Nishioka, R. S.; I. Yasumasu, A. Packard, H.; Bern and J. Z. Young. (1966). Nature of vesicles associated with the nervous system of cephalopods. *Z. Zellforsch*, 75:301-316.
- Nishioka, R., H. Bern and D. Golding (1970). Innervation of the cephalopod optic gland. In *Aspects of Neuroendocrinology*. W. Bargman & B. Schaer (eds). Springer, Berlin. p. 47-54.
- O'Dor, R. and M. J. Wells (1973). Yolk protein synthesis in the ovary of *Octopus vulgaris* and its control by the optic gland gonadotropin. *J. Exp. Biol.*, 59:665-674.
- Richard, A. (1966). La température, facteur essentiel de croissance pour le Céphalopode *Sepia officinalis*. *C. R. Acad. Sci., Paris. D.* 263:1138-1141.
- Richard, A. (1967a). Action de la température sur l'évolution génitale de *Sepia officinalis*. *C. R. Acad. Sci., Paris. D* 263:1998-2001.
- Richard, A. (1967b). Rôle de la photopériode dans le déterminisme de la maturation génitale femelle de Céphalopode *Sepia officinalis* L. *C. R. Acad. Sci., Paris. D* 265:1315-1318.. [
- Richard, A. (1970). Differentiation sexuelle de Céphalopodes en culture *in vitro*. *Ann. Biol.* IX(7, 9-10):409-415.
- Wells, M. and J. Wells (1959). Hormonal control of sexual maturity in *Octopus*. *J. Exp. Biol.* 36:1-33.
- Wells, M. and J. Wells (1969). Pituitary analogue in the *Octopus*. *Nature*, 222:293-294.
- Wells, M. and J. Wells (1975). Optic glands implants and their effects on the gonads of *Octopus*. *J. Exp. Biol.*, 62:579-588.
- Wells, M. J. (1960). Optic glands implants and the ovary of *Octopus*. *Symp. Zool. Soc. Lond.*, 2:87-107.
- Wodinsky, J. (1977). Hormonal inhibition of feeding in *Octopus* control by optic gland secretion. *Science*, 198:948-951.




Fig. 1. Dorsal view of the brain of *Sepioteuthis lessoniana* showing the optic gland (opt.gl.) associated with the optic lobe. The other brain structures are also shown (ventricle lobe:v.l., buccal ganglion: buc.g., palliovisceral ganglion: palv.g.).




Fig. 2. Transverse section of the brain of juvenile *S. lessoniana* showing the left optic gland (opt.gl.) in relation to optic lobe (opt.l) and optic fibres (opt.f.) and peduncle lobe (ped.l). Stained with Hx-E., 50X

A study on the optic glands of *Sepioteuthis lessoniana* from the Red Sea




Fig. 3. Transverse section of the brain of adult *S. lessoniana* showing the left optic gland (opt.gl.) in relation to optic lobe (opt.l) and optic tract fibres (opt.tr.f.), olfactory lobe (olf.l.) and peduncle lobe (ped.l.). Stained with Cajal Silver impregnation , Magn.100X




Fig. 4. Cells of optic gland in immature *S. lessoniana* showing its small size and blood vessels. Stained with Hx-E, 100X.




Fig. 5. Cells of optic gland in mature female *S. lessoniana* showing its large size cells with rounded nuclei and granulated cytoplasm. Stained with Hx-E, 200X (n: nucleus , nuc: nucleolus, s.gr.: secretory granules)




Fig. 6. Neurosecretory cells of optic gland in mature female *S. lessoniana* showing the stained granules scattered in the cytoplasm and the rounded nuclei and granulated cytoplasm. Stained with Paraldehyde fuchsin, 200X.

A study on the optic glands of *Sepioteuthis lessoniana* from the Red Sea




Fig. 7. Enlarged neurosecretory cell of optic gland in mature *S. lessoniana* showing the stained granules scattered in the cytoplasm and its axon. The nucleus is rounded with three nucleoli . Stained with Paraldehyde fuchsin, 200X. (ax.h.: axon helick, n: nucleus, s.gr.: secretory granules).

دراسة على الغدة البصرية للسبادج سيبوتيوسيس ميسونيانا من البحر الاحمر

وحيد محمود امام ، طارق على غريب
قسم علم الحيوان كلية العلوم جامعة عين شمس

تم في هذه الدراسة فحص الغدة البصرية لسبادج الشعاب المرجانية سيبوتيوسيس ميسونيانا من منطقة خليج السويس على البحر الاحمر. وتبين من الدراسة وجود غدتين صغيرتين يقع كل منها أعلى منطقة السرة من الخارج للفص البصرى عند اتصاله بالفص الساقى والفص الشمى. وأوضح التركيب الهيستولوجى ان هذه الغدة تتكون من خلايا كبيرة الحجم يصل قطرها الى 60 ميكرون ذات انوية دائرية (20 ميكرون) وتحتوى الخلايا سيتوبلازم محبب. وقد أوضحت الدراسة وجود حبيبات تصبغ بصبغة البرالدهيد فيكسين وهي تعتبر من الخلايا العصبية المفرزة وهي متصلة بالوعية الدموية. كما وجد ان هذه الغدة تتصل داخليا من خلال الياف العصب البصرى بالفصوص الظهرية الجانبية للمخ. وقد تم مناقشة وظيفة هذه الغدة فى عملية النضوج الجنسى للحيوان.